arXiv:cond-mat/0512486v1 [cond-mat.stat-mech] 20 Dec 2005

Collective dispersion relations for the 1D interacting two-component Bose and Fermi gases

M.T. Batchelor, M. Bortz, X.W. Guan and N. Oelkers

† Department of Theoretical Physics, RSPSE, and
Department of Mathematics, MSI
The Australian National University, Canberra ACT 0200, Australia

Abstract. We investigate the elementary excitations of charge and spin degrees for the 1D interacting two-component Bose and Fermi gases by means of the discrete Bethe ansatz equations. Analytic results in the limiting cases of strong and weak interactions are derived, where the Bosons are treated in the repulsive and the fermions in the strongly attractive regime. We confirm and complement results obtained previously from the Bethe ansatz equations in the thermodynamic limit.

PACS numbers: 03.75.Ss, 05.30.Fk, 71.10.Pm

1. Introduction

Recent achievements in trapping quantum gases of ultra-cold atoms have opened up many exciting possibilities for the experimental investigation of quantum effects in low-dimensional many-body systems. In particular, the experimental realization of the Tonks-Girardeau gas [1, 2, 3, 4, 5], collective spin wave excitations in the spinor Bose gas [6, 7, 8], the quasi 1D Fermi gas [9, 10] and the observation of atom matter waves [11] have shed light on understanding the quantum nature of 1D many-body systems. There has been a corresponding revival of interest in the exactly solved models of interacting Bose and Fermi gases, such as the Lieb-Liniger Bose gas [12], the Gaudin-Yang Fermi gas [13], the 1D Hubbard model [14, 15] and the BCS pairing model [16], due to the fact that their ground state and low lying excitations are accessible from exact Bethe ansatz solutions.

Quantum gases with multi-spin (pseudo-spin) states exhibit even richer quantum effects than their single component counterparts. The update facilities allow one to create multi-component Bose gases in low dimensions, e.g., two atomic hyperfine states can make up a pseudo-spin doublet [6, 7, 8]. The observation of collective dynamics of spin waves in the spinor Bose gas has stimulated theoretical attention on the 1D integrable spinor Bose gas [17, 18, 19, 20]. In this paper, we present a systematic way to derive dispersion relations of low-lying elementary excitations of the spinor Bose and Fermi gases via the discrete Bethe ansatz equations in the strong and weak coupling limits. This approach has some distinct advantages compared to the usual way of deriving the low-lying excitation spectrum: (i) it leads to explicit forms of the quasimomenta which are helpful in understanding the collective behaviour of low-lying excitations; (ii) it avoids the counting of quantum numbers involved in the integral equation formulation of the Bethe equations, and (iii) physical quantities such as the charge velocity, spin velocity, effective mass, Fermi momentum, finite-size corrections, ground state energy and excitation energies readily follow in the weak- and strong-coupling limits. Note that especially in weak coupling limit, those quantites are difficult to deal with in the framework of the Bethe ansatz equations in the thermodynamic limit (due to singular integral kernels in the linear integral equations).

To describe elementary excitations in terms of their dispersion relations, consider the general case of $\mathcal{M}$ branches (or rapidities) of excitations labeled by $\mu = 1, \ldots, M$. For example, $\mathcal{M} = 1$ for the spinless Bose gas (charge excitations only), $\mathcal{M} = 2$ for the spinor Bose and spin-$\frac{1}{2}$ Fermi gas (spin and charge excitations). Generally, the relativistic dispersion relation for each one of these excitations is written as

$$\varepsilon_\mu(p) = \sqrt{\Delta_\mu^2 + v_\mu^2 p^2} \tag{1}$$

where Δ_μ is the excitation gap and v_μ is the velocity. For vanishing gap $\Delta_\mu = 0$, the spectrum is linear, such that

$$v_\mu = \partial_p \varepsilon_\mu(p). \tag{2}$$

In this case, the underlying effective field theory is conformally invariant. Examples are the charge channel for spinless bosons with repulsive interaction [21], the charge channel for fermions and the spin channel for repulsive fermions. In the other extreme $\Delta_\mu \gg 1$, one ends up with

$$\varepsilon_\mu(p) = \Delta_\mu + \frac{v_\mu^2 p^2}{2\Delta_\mu} := \Delta_\mu + \frac{p^2}{2m_\mu^*} \tag{3}$$

which is the classical dispersion of free particles with an effective mass m_μ^*. It is known that this scenario is realized for spin excitations in the spinor Bose gas with repulsive interaction [22, 23]. This case has been analyzed recently [18, 20]. We will consider this model with our method in Section 2, thereby confirming quantitatively the following results. In the weak coupling limit, the effective mass of the spin wave is almost the same as the mass of one boson, while for strong coupling, it tends to the total mass of all bosons. On the other hand, charge excitations in this system are still gapless.

In Section 3, we will turn our attention to the intermediate regime with finite Δ_μ, which is encountered for spin excitations in the attractive spin-$\frac{1}{2}$ Fermi gas. In this system, the gap is an increasing function of c^2. The charge excitations again remain gapless. For the sake of comparison, we briefly recall the calculation of the velocities for the attractive spin-$\frac{1}{2}$ Fermi gas in the thermodynamic limit at the end of Section 3. We note that the method we propose to calculate $\varepsilon(p)$ in (1) can also be adapted to other models such as the 1D integrable Hubbard model [14, 15] and the mixed Bose-Fermi model [24, 25, 26, 27].

2. The spinor Bose gas

Before turning to the spinor Bose gas, we give an outline of the method we apply.

2.1. Outline of the method

In the framework of the Bethe ansatz solution, the energy eigenvalues E and momenta p of the two-component Bose and Fermi gases are given in terms of two sets of Bethe roots, $\{k_j\}_{j=1,\ldots,N}$ and $\{\lambda_\alpha\}_{\alpha=1,\ldots,M}$, corresponding to charge and spin degrees of freedom. Namely,

$$E = \sum_{j=1}^{N} k_j^2 \tag{4}$$

$$p = \sum_{j=1}^{N} k_j \tag{5}$$

where $\{k_j\}_{j=1,\ldots,N}$ and $\{\lambda_\alpha\}_{\alpha=1,\ldots,M}$ satisfy a set of $N+M$ coupled Bethe equations, given below. The ground state momentum p_0 and the ground state energy E_0 are given uniquely by the location and the number of roots; especially, $p_0 = 0$. Starting from this ground state configuration, low-lying excitations are constructed by changing both the location and the number of the roots, thereby obtaining the corresponding momentum p and energy E, both parametrized by the root sets. From these, one obtains the dispersion relation $E(p)$ and the excitation energy $\varepsilon(p) = E(p) - E_0$, which is analyzed according to Eq. (1). The essential point is that the roots can be obtained explicitly in the limits of weak and strong coupling. The thermodynamic limit, which is essential for Eq. (1) to be defined at all, is carried out at the very end.

2.2. Derivation of results

The Hamiltonian

$$H = -\frac{\hbar^2}{2m}\sum_{i=1}^{N}\frac{\partial^2}{\partial x_i^2} + g\sum_{1\le i<j\le N}\delta(x_i - x_j) \tag{6}$$

describes a quantum gas consisting of N particles with equal masses in one dimension, where the particles interact via a pairwise δ-function potential. According to the symmetry of the wave function, these may be bosons, fermions or a mixture of both. Here, we will consider a two-component Bose gas [18] constrained by periodic boundary conditions to a line of length L. The interaction is attractive for $c < 0$ and repulsive for $c > 0$. The mass is denoted by m and g is the coupling constant which is written in terms of the scattering strength $c = 2/a_{1\mathrm{D}}$ as $g = \hbar^2 c/m$. An effective 1D scattering length $a_{1\mathrm{D}}$ can be expressed through the 3D scattering length for the bosons confined in a quasi 1D geometry. For convenience in calculations, we prefer to set $\hbar = 2m = 1$, and thus to work with the parameter c. However, units may be restored at any time by using the dimensionless coupling constant

$$\gamma = mg/(\hbar^2 n) = c/n. \tag{7}$$

The energy is given by Eq. (4), where the quasimomenta k_j satisfy the Bethe equations (BE) [18, 28, 33]

$$\begin{aligned}\exp(\mathrm{i}k_jL) &= -\prod_{\ell=1}^{N}\frac{k_j - k_\ell + \mathrm{i}\,c}{k_j - k_\ell - \mathrm{i}\,c}\prod_{\mu=1}^{M}\frac{k_j - \lambda_\mu - \frac{1}{2}\mathrm{i}c}{k_j - \lambda_\mu + \frac{1}{2}\mathrm{i}c}\\ \prod_{\ell=1}^{N}\frac{\lambda_\alpha - k_\ell - \frac{1}{2}\mathrm{i}c}{\lambda_\alpha - k_\ell + \frac{1}{2}\mathrm{i}c} &= -\prod_{\beta=1}^{M}\frac{\lambda_\alpha - \lambda_\beta - \mathrm{i}\,c}{\lambda_\alpha - \lambda_\beta + \mathrm{i}\,c}.\end{aligned} \tag{8}$$

Here $j = 1, \ldots, N$ and $\alpha = 1, \ldots, M$, with M the number of spin-down bosons.

The ground state is the fully polarized state (total spin $S = N/2$) with $2S+1$-fold degeneracy due to $su(2)$ symmetry [17, 33]. This is achieved by choosing $M = 0$ in Eq. (8). So in the ground state, the BE reduce to those of the Lieb-Liniger Bose gas, which have been extensively studied via various methods [29, 30, 31, 32, 34]. From previous work [12, 31, 32, 34] on the Lieb-Liniger Bose gas, the ground state energy for weak coupling, namely $Lc \ll 1$, is given by

$$E_0 \approx N(N-1)c/L. \tag{9}$$

In the strong coupling limit, i.e., as the strength of the repulsive interaction tends to infinity, the bosons behave like impenetrable fermions and the system is known as the Tonks-Girardeau gas. In this regime, the ground state energy $E_0 \approx$

$\frac{\pi^2}{3L^2}\left(N^2-1\right)(1-4n/c)$ follows directly from the asymptotic solutions to the BE [32]. Due to conformal invariance, the finite-size corrections to the ground state energy in the thermodynamic limit for strong coupling are given by

$$E_0(L,N)-Le_0^{(\infty)}=-\frac{\pi C v_c}{6L}+o(1/L^2) \tag{10}$$

where $v_c=v_F\,(1-4n/c)$ with the Fermi velocity $v_F=2\pi n$ and the central charge $C=1$. We notice that in the zero coupling regime $c\to 0$ the finite-size corrections for the ground state energy do not meet the relation (10), because $v_c=2\sqrt{cn}$ [12] (cf. also the discussion after Eq. (13)). This is due to the fact that for $c=0$, the Bose gas has dispersion $\hbar^2k^2/(2m)$, as can be seen from the Hamiltonian (6). In that case the model is not conformally invariant (conformal invariance requires a *linear* dispersion). However, for any finite c, one identifies a finite Fermi momentum k_F. Then the dispersion relation can again be linearized around $\pm k_F$, yielding an effective bosonic theory which describes the density fluctuations with wave vectors $k\sim\pm k_F$ [21]. This theory is conformally invariant due to the linearized spectrum.

Spin charge separation is a typical phenomenon in interacting many-body systems. We first concentrate on the elementary excitations in the charge sector. These have linear dispersion, where the sound velocity is obtained from Eq. (2). In terms of the Bethe roots, these excitations are created by particle-hole pairs with respect to the ground state configuration of the $\{k_j\}_{j=1,\ldots,M}$. Let us first consider the weak-coupling limit, such that $k_i\propto\sqrt{c/L}$, $i=1,\ldots,N$ [32]. Then, to order c^2, the equations

$$\cos(k_iL)\approx 1-2\sum_{\ell=1}^{N}\frac{c^2}{(k_i-k_\ell)^2}-4\sum_{\ell=1}^{N}\sum_{\ell<\ell'=2}^{N}\frac{c}{(k_i-k_\ell)}\frac{c}{(k_i-k_{\ell'})} \tag{11}$$

$$\sin(k_iL)\approx\sum_{\ell=1}^{N}\frac{2c}{(k_i-k_\ell)} \tag{12}$$

hold, which determine the roots of the BE (8) asymptotically. The lowest possible excitations are obtained from the set of equations [32]

$$\begin{aligned} k_N &\approx p+\frac{2c}{L}\sum_{\ell=1}^{N-1}\frac{1}{k_N-k_\ell}\\ k_j &\approx \frac{2c}{L}\sum_{\ell=1,\ell\neq j}^{N}\frac{1}{k_j-k_\ell} \end{aligned} \tag{13}$$

where $j=1,\ldots,N-1$ and the momentum p parametrizes the excitation. Here, $p=2\ell\pi/L$ with ℓ integer; however, we will treat p as a variable in the following, which will take continuous values in the thermodynamic limit. Note that p is introduced such that Eq. (5) holds. The excited energy in the long wavelength limit is easily calculated from equations (13), i.e., $E(p)\approx N(N-1)c/L+pk_B+p^2$, where the Fermi boundary $k_B\approx 2\sqrt{cn}$ is obtained from the Hermite polynomial description [35]. Therefore $E(p)-E_0\approx 2\sqrt{cn}p=v_cp$. Here the charge velocity is given by $v_c=2\sqrt{nc}$, which coincides with the result obtained from (9) according to the formula $v_c=\sqrt{\frac{2L}{n}\left[\frac{\partial^2E}{\partial L^2}\right]}$

[12]. From the Bethe roots (13), the collective effect is that if one particle jumps out of the Dirac sea the rest also rearranges.

Similarly, for the strong coupling limit $Lc \gg 1$, if the largest quasimomentum k_N jumps out of the Dirac sea, asymptotic roots of the BE (8) can be obtained from a $1/(Lc)$ expansion as

$$\begin{aligned} k_N &\approx \left(\frac{(N-1)\pi}{L} + p + \frac{2p}{Lc}\right)\left(1 - \frac{2N}{Lc}\right) \\ k_j &\approx \left(\frac{2n_j\pi}{L} + \frac{2p}{Lc}\right)\left(1 - \frac{2N}{Lc}\right) \end{aligned} \tag{14}$$

where $n_j = \pm 1, \pm 3, \ldots, \pm(N-3)/2, -(N-1)/2$ if N is odd. It follows that $E(p) - E_0 \approx \frac{2(N-1)\pi}{L}\left(1 - \frac{2N}{Lc}\right)^2 p \approx v_c p$ with the charge velocity $v_c = \frac{2N\pi}{L}\left(1 - \frac{4N}{Lc}\right)$ and the ground state energy $E_0 \approx \frac{\pi^2}{3L^2}N(N^2-1)(1 - \frac{4N}{Lc})$. Note that as $c \to \infty$ the velocity v_c becomes the Fermi velocity of noninteracting spinless fermions.

We now focus on the spin excitation with regard to one spin flipping. This corresponds to introducing one λ-solution into Eqs. (8), which we call λ_1. In the long wavelength limit a quadratic dispersion of spin wave excitations above the ferromagnetic ground state is obtained [22, 23, 20]

$$E(p) - E_0 \approx p^2/2m^* \tag{15}$$

where m^* is an effective mass. This is the case indicated in Eq. (3). Now the small-coupling expansion of the BE (8) yields

$$k_j \approx \frac{2c}{L}\sum_{\ell=1}^{N}\frac{1}{k_j - k_\ell} + \frac{c}{L\lambda_1} + \frac{c}{L\lambda_1^2}k_j \tag{16}$$

for $j = 1, \ldots, N$. From Eq. (5), one obtains $\lambda_1 \approx \frac{Nc}{Lp}$, which in the long-wavelength limit diverges as $1/p$, so that $\lambda_1 \gg k_B$. We now calculate the energy by inserting Eq. (16) into Eq. (4),

$$E(p) = \sum_{j=1}^{N} k_j^2 \approx \left[\frac{c}{L}N(N-1) + \frac{c}{L\lambda_1}p\right]\left(1 + \frac{c}{L\lambda_1^2}\right) = E_0 + p^2 \tag{17}$$

which indicates $m/m^* \approx 1$, where m is the boson mass. In the above equation E_0 is given by Eq. (9). We notice that higher order corrections to the ground state energy are very hard to derive from our expansion ansatz. The next correction to the ground state energy of the Lieb-Liniger Bose gas is known to be given by $E_0 \approx \frac{N^2c}{L}\left(1 - \frac{4\sqrt{c/n}}{3\pi}\right)$ [31]. Thus from the relation (17) we observe that

$$E(p) \approx E_0\left(1 + \frac{c}{L\lambda_1^2}\right). \tag{18}$$

This suggests the effective mass $m/m^* \approx 1 - \frac{4\sqrt{c/n}}{3\pi}$.

The divergence $\lambda_1 \sim 1/p$ continues to hold in the strong coupling regime $c \gg 1$. In this regime, the consistency conditions derived from the BE (8) in the $1/(Lc)$ expansion are

$$\sin k_j L \approx -2\sum_{\ell=1}^{N}\frac{k_j-k_\ell}{c}+\frac{c}{\lambda_1}\left(1+\frac{k_j}{\lambda_1}\right). \tag{19}$$

Here for convenience we consider N as an odd number. Then

$$k_j \approx \left(\frac{2\pi n_j}{L}+\frac{2p}{Lc}+\frac{c}{L\lambda_1}\right)\left(1-\frac{2N}{Lc}+\frac{c}{L\lambda_1^2}\right) \tag{20}$$

where $n_j = \pm 1, \pm 3, \ldots, \pm(N-1)/2$. From Eq. (5), we find $\lambda_1 \approx Nc/Lp \gg c \gg k_j$. After some algebra we obtain

$$\begin{aligned} E(p) = \sum_{j=1}^{N} k_j^2 &\approx \frac{\pi^2}{3L^2}N(N^2-1)\left(1-\frac{4N}{Lc}\right) \\ &+ p^2\left(\frac{1}{N}+\frac{2\pi^2 N}{3Lc}\left(1+\frac{2N}{Lc}\right)+\frac{4}{Lc}\right)\left(1-\frac{4N}{Lc}\right) \end{aligned} \tag{21}$$

which suggests the effective mass

$$\frac{m}{m^*} \approx \frac{1}{N}+\frac{2\pi^2 n}{3c}\left(1-\frac{2n}{c}\right). \tag{22}$$

The leading correction term coincides with the earlier result [20]. The next correction term is relevant for finite c. In the strong coupling limit, i.e., $c \to \infty$, all the bosons become indistinguishable thus behaving like hardcore bosons with Fermi pressure-like kinetic energy. The effective mass takes the maximum value $m^* = Nm$, meaning that by moving one boson with down spin, one has to move all the particles with up spins. In this way the spin velocity is very small, as if the spin-flip is frozen [20].

3. The Fermi gas

The spin-$\frac{1}{2}$ Fermi gas with equal number of spin-up and spin-down fermions described by the Hamiltonian (6) has been extensively studied in the context of the BCS-BEC crossover [36, 37, 38, 39]. Very recently, a comprehensive study of the model with arbitrary polarization was undertaken [40]. In order to compare different collective dynamics of elementary excitations in the charge and spin degrees with those of the spinor Bose gas, we derive the charge and spin velocities for the strongly attractive Fermi gas. This is done from an expansion of the discrete BE, similarly to the previous section. At the end of this section, we will compare the results with those obtained from the BE in the thermodynamic limit.

3.1. Velocities from the discrete Bethe ansatz equations

In the strongly attractive regime the bound states behave like bosonic tightly bound molecular dimers. The ground state consists of two-body bound states spread out about the origin in pseudomomentum space due to Fermi pressure, i.e., with $k_j = \lambda_j + \frac{1}{2}\mathrm{i}c$, $k_{-j} = \lambda_j - \frac{1}{2}\mathrm{i}c$ for $j = 1, \ldots, N/2$, where the parameters $\{\lambda_j\}_{j=1,\ldots,M}$ satisfy the BE [13, 30]

$$\mathrm{e}^{\mathrm{i}\,2\lambda_j L} = -\prod_{\ell=1}^{M} \frac{\lambda_j - \lambda_\ell + \mathrm{i}c}{\lambda_j - \lambda_\ell - \mathrm{i}c}, \qquad M = N/2 \tag{23}$$

for $j = 1, \ldots, M$. The lowest charge excitation comes from the configuration in which a bound state with the largest momentum $k_{\max} = \frac{(M-1)\pi}{L}(1 - \frac{M}{Lc})$, which is at the edge of the Dirac sea in quasimomentum space, jumps out of the Dirac sea. This leads to a bound state density fluctuation. The total real momentum changes from zero to p. Taking M even, we may obtain the roots of the BE (23) as

$$\begin{aligned} \lambda_M &\approx \left[\frac{(M-1)\pi}{2L} + \frac{p}{2} + \frac{p}{2Lc}\right]\left(1 - \frac{M}{Lc}\right) \\ \lambda_j &\approx \left[\frac{2n_j - 1}{2L} + \frac{p}{2Lc}\right]\left(1 - \frac{M}{Lc}\right) \end{aligned} \tag{24}$$

where $j = 1, \ldots, M-1$ and $n_j = -(M-2)/2, -(M-4)/2, \ldots, (M-2)/2$. It follows that

$$E(p) \approx -\frac{1}{4}c^2 N + \frac{\pi^2}{48L^2}N(N^2-2)\left(1 - \frac{2M}{Lc}\right) + \frac{(M-1)\pi}{L}\left(1 - \frac{2M}{Lc}\right)p. \tag{25}$$

It is seen that to this order $E(p) - E_0 = v_c\, p$ with charge velocity

$$v_c = \frac{n\pi}{2}\left(1 - \frac{n}{c}\right) \tag{26}$$

which coincides with the result of Ref. [36] and with the result derived from the ground state energy E_0 [36, 37, 39, 40],

$$E_0 = -\frac{1}{4}c^2 N + \frac{\pi^2}{48L^2}N(N^2-2)\left(1 - \frac{2M}{Lc}\right). \tag{27}$$

Again we may find from the ground state energy that the finite-size corrections in the strongly repulsive limit for the Fermi gas also meet the relation (10) with central charge $C = 1$. The spin-channel does not contribute here, because it is gapped (cf. below).

We now focus on the spin velocity which involves breaking one bound state over the ground state in order to get spin flipping. The ground state is a spin singlet. The lowest spin excitation is incurred by a spin triplet excitation with total spin $M^z = 1$. In such a configuration the BE become [30]

$$\begin{aligned} \mathrm{e}^{\mathrm{i}\,2\lambda_j L} &= -\prod_{\ell=1}^{M-1} \frac{\lambda_j - \lambda_\ell + \mathrm{i}c}{\lambda_j - \lambda_\ell - \mathrm{i}c} \prod_{\mu=1}^{2} \frac{\lambda_j - k_\mu + \frac{1}{2}\mathrm{i}c}{\lambda_j - k_\mu - \frac{1}{2}\mathrm{i}c} \\ \mathrm{e}^{\mathrm{i}\,k_\mu L} &= -\prod_{\ell=1}^{M-1} \frac{k_\mu - \lambda_\ell + \frac{1}{2}\mathrm{i}c}{k_\mu - \lambda_\ell - \frac{1}{2}\mathrm{i}c} \end{aligned} \tag{28}$$

with $j = 1, \ldots M-1$ and $\mu = 1, 2$. Withought loss of generality, we assume M is even. The BE (28) admit the asymptotic roots

$$k_1 \approx \left(\frac{n_+\pi}{L} + \frac{4}{Lc}\sum_{j=1}^{M-1}\lambda_j\right)\left(1 - \frac{4(M-1)}{Lc}\right)$$

$$
\begin{aligned}
k_2 &\approx \left(\frac{n_-\pi}{L} + \frac{4}{Lc}\sum_{j=1}^{M-1}\lambda_j\right)\left(1 - \frac{4(M-1)}{Lc}\right) \\
\lambda_j &\approx \left(\frac{n_j\pi}{L} + \frac{1}{Lc}\sum_{j=1}^{M-1}\lambda_j + \frac{2(k_1+k_2)}{Lc}\right)\left(1 - \frac{M-3}{Lc}\right).
\end{aligned} \tag{29}
$$

In the above equations $n_j = 0, \pm1, \cdots, \pm(M-2)/2$ and $n_\pm$ are odd numbers to be determined. The number of bound states is now $M-1$. This leads to a different parity for the BE (28). We can see this directly from the roots (29) where all bound states shift by nearly $\pi/2$ from their original positions in the ground state configuration. The unpaired two fermions are scattered away. Summing up both sides of the third equation in Eq. (29) gives the relation $\sum_{j=1}^{M-1}\lambda_j \approx \frac{2(M-1)}{Lc}(k_1+k_2)$. This relation implies that $\sum_{j=1}^{M-1}\lambda_j$ is negligible in comparison with k_1+k_2 in the strong coupling regime. Further, from Eq. (5) we have

$$
k_1 + k_2 \approx p\left(1 - \frac{4(M-1)}{Lc}\right) \tag{30}
$$

if we consider $n_- = -n_+$ for minimizing the energy. Using the BE in (28) gives another relation

$$
k_1 - k_2 \approx \frac{2n_+\pi}{L}\left(1 - \frac{4(M-1)}{Lc}\right). \tag{31}
$$

Thus we find all roots to be given by

$$
\begin{aligned}
\lambda_j &\approx \left(\frac{n_j\pi}{L} + \frac{2p}{Lc}(1 - \frac{4(M-1)}{Lc})\right)\left(1 - \frac{M-3}{Lc}\right) \\
k_{1,2} &\approx \left(\pm\frac{n_+\pi}{L} + \frac{p}{2}\right)\left(1 - \frac{4(M-1)}{Lc}\right).
\end{aligned} \tag{32}
$$

Here $n_j = 0, \pm1, \ldots, \pm(M-2)/2$. The quantum numbers $n_\pm$ for the unpaired fermions should be chosen to be as small as possible, i.e., ±1, in order to minimize the energy [30]. Actually, $n_\pm$ may take any two odd integer values, which correspond to states with energy higher than the ground state energy. After some algebra, we obtain the excitation energy

$$
\begin{aligned}
E(p) \approx & -2(M-1)\frac{c^2}{4} + \frac{\pi^2}{48L^2}\left(1 - \frac{2(M-3)}{Lc}\right)N(N^2-4) \\
& + \frac{1}{2}p^2\left(1 - \frac{8(M-1)}{Lc}\right) + O\left(\frac{n_+^2}{L^2}, \frac{M^2}{L^2}\right).
\end{aligned} \tag{33}
$$

The choice of $n_\pm$ affects the excitation energy only to the indicated order. From Eq. (33) we deduce the dispersion relation $\varepsilon(p) = E(p) - E_0 = \Delta + \frac{1}{2}p^2\left(1 - \frac{8M}{Lc}\right)$, where the spin excitation gap to leading order in c is $\Delta \approx c^2/2$ and E_0 is given by (27). On the other hand, by comparing with Eq. (1), the spin velocity follows as

$$
v_s = \sqrt{\Delta}\left(1 - \frac{4M}{Lc}\right) = \frac{|c|}{\sqrt{2}}\left(1 - \frac{2n}{c}\right) \tag{34}
$$

which is divergent due to the gapped spin triplet excitation [41, 40, 42]. This result has been obtained previously in Ref. [36].

3.2. Velocities from the Bethe ansatz equations in the thermodynamic limit

Spin and charge velocites for the spin-$\frac{1}{2}$ Fermi gas in the attractive regime have been calculated [36, 40], using the BE in the thermodynamic limit. Within this approach, the ideas sketched in Section 2.1 are applied to the log form of Eqs. (8), thereby converting products to sums. By introducing densities, these can in turn be converted to integrals. One ends up with linear integral equations which can be solved numerically in order to determine v_c and v_s. The advantage of this approach is that it is valid for arbitrary couplings. In Fig. 1, v_c and v_s are depicted from a numerical solution of those integral equations for attractive fermions. Furthermore, the results (26), (34), are confirmed.



Figure 1. a) Charge and spin velocity divided by $v_F = \pi n$ for the attractive regime, dependent on γ. The dashed lines are the small γ approximations, $v_{c,s}/v_F = 1 \pm \gamma/\pi^2$. b) The corrections to $v_{c,s}$ in the strongly attractive limit, namely $-\gamma(2v_c/v_F - 1)$ and $(\pi\sqrt{2}v_s/(v_F\gamma) + 1)\gamma/2$. According to Eqs. (26), (34) and (7), these expressions approach 1 in the limit $\gamma \to -\infty$, which is confirmed numerically.

4. Concluding remarks

In conclusion, we have derived explicitly the dispersion relations for elementary charge and spin excitations for the two-component Bose and Fermi gases in the long wavelength limit. Analytic results have been found for the charge and spin velocities and for the effective mass connected with spin wave excitations in the repulsive spinor Bose gas. These quantities reveal significant insight into the nature of the collective hydrodynamics of the Bose and Fermi gases. In particular, physical quantities such as the charge velocity, spin velocity, effective mass, Fermi momentum, finite-size corrections, ground state and excitation energies, are obtained via an analysis of the discrete Bethe ansatz solutions, as opposed to using the continuous integral equation approach. As such we believe that it can be easily adapted to a number of related models.

Acknowledgments

This work has been supported by the Australian Research Council and the German Science Foundation under grant number BO 2538/1-1. We thank D. M. Gangardt, C. Lee, S. Sergeev and H.-Q. Zhou for helpful discussions. XWG also thanks the Department of Mathematics, University of Science and Technology of China, for kind hospitality.

References

[1] Greiner M *et al* 2002 *Nature* **415** 39
[2] Paredes B *et al* 2004 *Nature* **429** 277
[3] Pollet L, Rombouts S M A and Denteneer P J H 2004 *Phys. Rev. Lett.* **93** 210401
[4] Kinoshita T, Wenger T and Weiss D S 2004 *Science* **305** 1125
[5] Moritz H, Stöferle T, Köhl M and Esslinger T 2003 *Phys. Rev. Lett.* **91** 250402
[6] Lewandowski H J, Harber D M, Whitaker D L and Cornell E A 2002 *Phys. Rev. Lett.* **88** 070403
[7] McGuirk J M et al 2002 *Phys. Rev. Lett.* **89** 090402
[8] McGuirk J M, Harber D M, Lewandowski H J and Cornell E A 2003 *Phys. Rev. Lett.* **91** 150402
[9] Moritz H et al 2005 *Phys. Rev. Lett.* **94** 210401
[10] Partridge G B, Li W, Kamar R I, Liao Y and Hulet R G cond-mat/0511752
[11] Perreault J D and Cronin A D 2005 *Phys. Rev. Lett.* **95** 133201
[12] Lieb E H and Liniger W 1963 *Phys. Rev.* **130** 1605
Lieb E H 1963 *Phys. Rev.* **130** 1616
[13] Gaudin M 1967 *Phys. Lett. A* **24** 1312
Yang C N 1967 *Phys. Rev. Lett.* **19** 1312
[14] Recati A *et al* 2003 *Phys. Rev. Lett.* **90** 020401
Liu X J Drummond D and Hu H 2005 *Phys. Rev. Lett.* **94** 136406
[15] Amico L, Mastellone A and Osterloh A cond-mat/0509554
[16] Ortiz G and Dukelsky J 2005 arXiv:cond-mat/0503664
[17] Eisenberg E and Lieb E H 2002 *Phys. Rev. Lett.* **89** 220403
[18] Li Y Q, Gu S J, Ying Z J and Eckern U 2003 *Europhys. Lett.* **61** 368
[19] Ieda J, Miyakawa T and Wadati M 2004 *Phys. Rev. Lett.* **93** 194102
[20] Fuchs J N, Gangardt D M, Keilmann T and Shlyapnikov G V 2005 *Phys. Rev. Lett.* **95** 150402
[21] Haldane F D M 1981 *Phys. Rev. Lett.* **47** 1840
[22] Halperin B I and Hohenberg P C 1969 *Phys. Rev.* **188** 898
[23] Halperin B I 1975 *Phys. Rev. B* **11** 178
[24] Lai C K and Yang C N 1971 *Phys. Rev. A* **3** 393
[25] Imambekov A and Demler E 2005 arXiv:cond-mat/0505632
—— arXiv:cond-mat/0510801
[26] Batchelor M T, Bortz M, Guan X W and Oelkers N 2005 *Phys. Rev. A* **72** 061603(R)
[27] Frahm H and Palacios G 2005 *Phys. Rev. A* **3** **72** 061604(R)
[28] Gu S J, Li Y Q, Ying Z J and Zhao X A 2002 *Int. J. Mod. Phys. B* **16** 2137

[29] Korepin V E, Izergin A G and Bogoliubov N M 1993 *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge)

[30] Takahashi M 1999 *Thermodynamics of One-Dimensional Solvable Models* (Cambridge University Press, Cambridge)

[31] Wadati M 2002 *J. Phys. Soc. Jpn* **71** 2657

[32] Batchelor M T, Guan X W and McGuire J B 2004 *J. Phys. A: Math. Gen.* **37** L497
Batchelor M T, Guan X W, Dunning C and Links J 2005 *J. Phys. Soc. Jpn* **74** 53 Suppl.

[33] Oelkers N, Batchelor M T, Bortz M and Guan X-W cond-mat/0511694

[34] Sakmann K, Streltsov A I, Alon O and Cederbaum L S 2005 *Phys. Rev. A* **72** 033613

[35] Hahn W 1934 *Bericht über die Nullstellen der Laguerreschen und der Hermiteschen Polynome* , Jahresbericht der deutschen Mathematiker Vereinigung **44** 215

[36] Fuchs J N, Recati A and Zwerger W 2004 *Phys. Rev. Lett.* **93** 090408

[37] Astrakharchik G E, Blume D, Giorgini S and Pitaevskii L P *Phys. Rev. Lett.* **93** 050402

[38] Tokatly I V *Phys. Rev. Lett.* **93** 090405

[39] Iida T and Wadati M 2005 *J. Phys. Soc. Jpn* **74** 1724

[40] Batchelor M T, Bortz M, Guan X W and Oelkers N 2005 arXiv:cond-mat/0506264

[41] Krivnov V Y and Ovchinnikov A A 1975 *Sov. Phys. JETP* **40** 781

[42] Cazallilla M A, Ho A F and Giamarchi T 2005 *Phys. Rev. Lett.* **95** 226402